arXiv:1008.2113v1 [gr-qc] 12 Aug 2010

# Electromagnetic field with induced massive term: Case with spinor field

Yu.P. Rybakov, G.N. Shikin and Yu.A. Popov

*Department of Theoretical Physics*
*Peoples' Friendship University of Russia*
*117198 Moscow, Russia*[*]

Bijan Saha

*Laboratory of Information Technologies*
*Joint Institute for Nuclear Research, Dubna*
*141980 Dubna, Moscow region, Russia*[†]

(Dated: August 13, 2010)

We consider an interacting system of spinor and electromagnetic field, explicitly depending on the electromagnetic potentials, i.e., interaction with broken gauge invariance. The Lagrangian for interaction is chosen in such a way that the electromagnetic field equation acquires an additional term, which in some cases is proportional to the vector potential of the electromagnetic field. This equation can be interpreted as the equation of motion of photon with induced non-trivial rest-mass. This system of interacting spinor and scalar fields is considered within the scope of Bianchi type-I (BI) cosmological model. It is shown that, as a result of interaction the electromagnetic field vanishes at $t \to \infty$ and the isotropization process of the expansion takes place.

PACS numbers: 03.65.Pm and 04.20.Ha
Keywords: electromagnetic field, spinor field, Bianchi type I (BI) model, photon mass

## I. INTRODUCTION

Though the triumphs of Maxwellian electromagnetism and quantum electrodynamics set the rest mass of photon to be trivial, the hypothesis of possible nontrivial photon mass has long been discussed in the literature [1–4]. The modern experimental data do not contradict this hypothesis [5–13]. So it is interesting to consider some additional arguments for or against this hypothesis. As one of such arguments can serve experimental data of modern observational cosmology, which witnesses the isotropy of the Universe. It is interesting to combine this fact with the description of matter by means of system of interacting fields including the electromagnetic one. In a recent paper [14] we considered one of the simplest systems comprising with mass-less scalar and electromagnetic fields and study the influence of such interaction on the expansion of the Universe in the asymptotic region. In that paper it was shown that if one consider only electromagnetic field, the two of the three spatial components vector potential are either constant or zero and the space-time in this case does not allow isotropization.

---

[*] Electronic address: soliton4@mail.ru
[†] Electronic address: bijan@jinr.ru; URL: `http://bijansaha.narod.ru`

In the recent years system with nonlinear spinor field was extensively studied in different cosmological models and it was shown that the nonlinear spinor field plays very important role in (i) isotropization of initially anisotropic space-time, (ii) formation of singularity free cosmological solutions, and (iii) explaining late-time acceleration [15–19]. In connection with this in this paper we consider an system of spinor and electromagnetic fields within the scope of a Bianchi type-I cosmological model and examine the influence of such interaction on the expansion of the Universe in the asymptotic region.

## II. BASIC EQUATIONS AND THEIR GENERAL SOLUTIONS

We choose the Lagrangian of the interaction electromagnetic and spinor fields within the framework of a BI cosmological gravitational field in the form

$$\mathscr{L} = \frac{R}{2\kappa} - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{i}{2}\left[\bar{\psi}\gamma^{\mu}\nabla_{\mu}\psi - \nabla_{\mu}\bar{\psi}\gamma^{\mu}\psi\right] - m\bar{\psi}\psi + \frac{1}{2}K(I)\mathscr{D}(S), \tag{2.1}$$

with $I = A_{\lambda}A^{\lambda}$ and $S = \bar{\psi}\psi$. We consider the BI metric in the form

$$ds^2 = e^{2\alpha}dt^2 - e^{2\beta_1}dx^2 - e^{2\beta_2}dy^2 - e^{2\beta_3}dz^2. \tag{2.2}$$

The metric functions $\alpha, \beta_1, \beta_2, \beta_3$ depend on $t$ only and obey the coordinate condition

$$\alpha = \beta_1 + \beta_2 + \beta_3. \tag{2.3}$$

Written in the form

$$R_{\mu}^{\nu} = -\kappa\Big(T_{\mu}^{\nu} - \frac{1}{2}\delta_{\mu}^{\nu}T\Big), \tag{2.4}$$

the Einstein equations corresponding to the metric (2.2) in account of (2.3) read

$$e^{-2\alpha}\Big(\ddot{\alpha} - \dot{\alpha}^2 + \dot{\beta}_1^2 + \dot{\beta}_2^2 + +\dot{\beta}_3^2\Big) \;=\; -\kappa\Big(T_0^0 - \frac{1}{2}T\Big), \tag{2.5a}$$

$$e^{-2\alpha}\ddot{\beta}_1 \;=\; -\kappa\Big(T_1^1 - \frac{1}{2}T\Big), \tag{2.5b}$$

$$e^{-2\alpha}\ddot{\beta}_2 \;=\; -\kappa\Big(T_2^2 - \frac{1}{2}T\Big), \tag{2.5c}$$

$$e^{-2\alpha}\ddot{\beta}_3 \;=\; -\kappa\Big(T_3^3 - \frac{1}{2}T\Big), \tag{2.5d}$$

where over dot means differentiation with respect to $t$ and $T_{\nu}^{\mu}$ is the energy-momentum tensor of the material field.

Variation of (2.1) with respect to electromagnetic field gives

$$\frac{1}{\sqrt{-g}}\frac{\partial}{\partial x^{\nu}}\Big(\sqrt{-g}F^{\mu\nu}\Big) - \mathscr{D}(S)\frac{dK}{dI}A^{\mu} = 0. \tag{2.6}$$

The spinor field equation corresponding to the metric (2.1) has the form

$$i\gamma^{\mu}\nabla_{\mu}\psi - m\psi + \frac{1}{2}K(I)\frac{d\mathscr{D}}{dS}\psi \;=\; 0, \tag{2.7a}$$

$$i\nabla_{\mu}\bar{\psi}\gamma^{\mu} + m\bar{\psi} - \frac{1}{2}K(I)\frac{d\mathscr{D}}{dS}\bar{\psi} \;=\; 0. \tag{2.7b}$$

The energy-momentum tensor of the interacting matters fields has the form

$$\begin{aligned} T_\mu^\nu &= -F_{\mu\eta}F^{\nu\eta} + \frac{i}{4}g^{\nu\rho}\Big(\bar\psi\gamma_\rho\nabla_\mu\psi + \bar\psi\gamma_\mu\nabla_\rho\psi - \nabla_\mu\bar\psi\gamma_\rho\psi - \nabla_\rho\bar\psi\gamma_\mu\psi\Big) \\ &+ \mathscr{D}(S)\frac{dK}{dI}A_\mu A^\nu - \delta_\mu^\nu\mathscr{L}. \end{aligned} \tag{2.8}$$

We consider the case when the electromagnetic and scalar fields are the functions of $t$ only. Taking this in mind we choose the vector potential in the following way:

$$A_\mu = \big(0, A_1(t), A_2(t), A_3(t)\big). \tag{2.9}$$

In this case the electromagnetic field tensor $F^{\mu\nu}$ has only three non-vanishing components, namely

$$F_{01} = \dot A_1, \quad F_{02} = \dot A_2, \quad F_{03} = \dot A_3. \tag{2.10}$$

On account of (2.9) and (2.10) we now have

$$I = A_\lambda A^\lambda = -A_1^2e^{-2\beta_1} - A_2^2e^{-2\beta_2} - A_3^2e^{-2\beta_3}, \tag{2.11}$$

$$F_{\mu\nu}F^{\mu\nu} = -2e^{-2\alpha}\big(\dot A_1^2e^{-2\beta_1} + \dot A_2^2e^{-2\beta_2} + \dot A_3^2e^{-2\beta_3}\big). \tag{2.12}$$

For electromagnetic field from (2.6) we find

$$\frac{d}{dt}\Big(\dot A_1e^{-2\beta_1}\Big) + \mathscr{D}(S)\frac{dK}{dI}e^{2\alpha-2\beta_1}A_1 = 0, \tag{2.13a}$$

$$\frac{d}{dt}\Big(\dot A_2e^{-2\beta_2}\Big) + \mathscr{D}(S)\frac{dK}{dI}e^{2\alpha-2\beta_2}A_2 = 0, \tag{2.13b}$$

$$\frac{d}{dt}\Big(\dot A_3e^{-2\beta_3}\Big) + \mathscr{D}(S)\frac{dK}{dI}e^{2\alpha-2\beta_3}A_3 = 0. \tag{2.13c}$$

Let us now go back to the spinor field equation. The spinor field equation for the metric (2.2) takes the form

$$ie^{-\alpha}\bar\gamma_0\Big(\partial_t + \frac{\dot\alpha}{2}\Big)\psi - m\psi + \frac{1}{2}K(I)\frac{d\mathscr{D}}{dS}\psi = 0, \tag{2.14}$$

where $\bar\gamma_\mu$ is the Dirac matrices for flat space-time. Equation (2.14) can be rewritten as

$$\dot\psi_a + \frac{\dot\alpha}{2}\psi_a + ie^\alpha\Big(m - \frac{1}{2}K(I)\frac{d\mathscr{D}}{dS}\Big)\psi_a = 0, \quad a = 1, 2, \tag{2.15a}$$

$$\dot\psi_b + \frac{\dot\alpha}{2}\psi_b - ie^\alpha\Big(m - \frac{1}{2}K(I)\frac{d\mathscr{D}}{dS}\Big)\psi_a = 0, \quad b = 3, 4. \tag{2.15b}$$

Solving (2.15) we find the following components of the spinor fields

$$\psi_a = C_a\exp\Big[-\frac{\alpha}{2} - i\int\Big(m - \frac{1}{2}K(I)\frac{d\mathscr{D}}{dS}\Big)e^\alpha dt\Big], \quad C_a = \text{const.} \tag{2.16a}$$

$$\psi_b = C_b\exp\Big[-\frac{\alpha}{2} + i\int\Big(m - \frac{1}{2}K(I)\frac{d\mathscr{D}}{dS}\Big)e^\alpha dt\Big], \quad C_b = \text{const.} \tag{2.16b}$$

From (2.16) we also find

$$S = S_0e^{-\alpha}. \tag{2.17}$$

Finally, let us solve the Einstein equations. In doing so, let us first write the non trivial components of the energy momentum tensor of material fields. In view of (2.15) from (2.8) we find

$$
\begin{aligned}
T_0^0 &= mS - \frac{1}{2}K(I)\mathscr{D}(S) + \frac{1}{2}\Big(\dot{A}_1^2 e^{-2\beta_1} + \dot{A}_2^2 e^{-2\beta_2} + \dot{A}_3^2 e^{-2\beta_3}\Big)e^{-2\alpha}, && \text{(2.18a)}\\
T_1^1 &= \frac{1}{2}K(I)\Big(\frac{d\mathscr{D}}{dS}S - \mathscr{D}(S)\Big) + \frac{1}{2}\Big(\dot{A}_1^2 e^{-2\beta_1} - \dot{A}_2^2 e^{-2\beta_2} - \dot{A}_3^2 e^{-2\beta_3}\Big)e^{-2\alpha}\\
&- \mathscr{D}\frac{dK}{dI}A_1^2 e^{-2\beta_1}, && \text{(2.18b)}\\
T_2^2 &= \frac{1}{2}K(I)\Big(\frac{d\mathscr{D}}{dS}S - \mathscr{D}(S)\Big) + \frac{1}{2}\Big(\dot{A}_2^2 e^{-2\beta_2} - \dot{A}_3^2 e^{-2\beta_3} - \dot{A}_1^2 e^{-2\beta_1}\Big)e^{-2\alpha}\\
&- \mathscr{D}\frac{dK}{dI}A_2^2 e^{-2\beta_2}, && \text{(2.18c)}\\
T_3^3 &= \frac{1}{2}K(I)\Big(\frac{d\mathscr{D}}{dS}S - \mathscr{D}(S)\Big) + \frac{1}{2}\Big(\dot{A}_3^2 e^{-2\beta_3} - \dot{A}_1^2 e^{-2\beta_1} - \dot{A}_2^2 e^{-2\beta_2}\Big)e^{-2\alpha}\\
&- \mathscr{D}\frac{dK}{dI}A_3^2 e^{-2\beta_3}, && \text{(2.18d)}\\
T_1^2 &= \Big(\dot{A}_1\dot{A}_2 e^{-2\alpha} - \mathscr{D}\frac{dK}{dI}A_1A_2\Big)e^{-2\beta_1}. && \text{(2.18e)}\\
T_2^3 &= \Big(\dot{A}_2\dot{A}_3 e^{-2\alpha} - \mathscr{D}\frac{dK}{dI}A_2A_3\Big)e^{-2\beta_2}. && \text{(2.18f)}\\
T_3^1 &= \Big(\dot{A}_3\dot{A}_1 e^{-2\alpha} - \mathscr{D}\frac{dK}{dI}A_3A_1\Big)e^{-2\beta_3}. && \text{(2.18g)}
\end{aligned}
$$

From (2.18) one also finds

$$
T = mS + \frac{3}{2}K(I)\frac{d\mathscr{D}}{dS}S - 2K(I)\mathscr{D}(S) - \mathscr{D}\frac{dK}{dI}\Big(A_1^2 e^{-2\beta_1} + A_2^2 e^{-2\beta_2} + A_3^2 e^{-2\beta_3}\Big). \tag{2.19}
$$

In view of (2.18) and (2.19) system of Einstein equations now takes the form

$$
\begin{aligned}
\ddot{\alpha} - \dot{\alpha}^2 + \dot{\beta}_1^2 + \dot{\beta}_2^2 + \dot{\beta}_3^2 &= -\frac{\kappa e^{2\alpha}}{2}\Big[\Big(mS + K\mathscr{D} - \frac{3}{2}K\frac{d\mathscr{D}}{dS}S\Big) + \Big(\dot{A}_1^2 e^{-2\alpha} + \mathscr{D}\frac{dK}{dI}A_1^2\Big)e^{-2\beta_1}\\
&+ \Big(\dot{A}_2^2 e^{-2\alpha} + \mathscr{D}\frac{dK}{dI}A_2^2\Big)e^{-2\beta_2} + \Big(\dot{A}_3^2 e^{-2\alpha} + \mathscr{D}\frac{dK}{dI}A_3^2\Big)e^{-2\beta_3}\Big], && \text{(2.20a)}\\
\ddot{\beta}_1 &= -\frac{\kappa e^{2\alpha}}{2}\Big[-\Big(mS + \frac{1}{2}K\frac{d\mathscr{D}}{dS}S - K\mathscr{D}\Big) + \Big(\dot{A}_1^2 e^{-2\alpha} - \mathscr{D}\frac{dK}{dI}A_1^2\Big)e^{-2\beta_1}\\
&- \Big(\dot{A}_2^2 e^{-2\alpha} - \mathscr{D}\frac{dK}{dI}A_2^2\Big)e^{-2\beta_2} - \Big(\dot{A}_3^2 e^{-2\alpha} - \mathscr{D}\frac{dK}{dI}A_3^2\Big)e^{-2\beta_3}\Big], && \text{(2.20b)}\\
\ddot{\beta}_2 &= -\frac{\kappa e^{2\alpha}}{2}\Big[-\Big(mS + \frac{1}{2}K\frac{d\mathscr{D}}{dS}S - K\mathscr{D}\Big) + \Big(\dot{A}_2^2 e^{-2\alpha} - \mathscr{D}\frac{dK}{dI}A_2^2\Big)e^{-2\beta_2}\\
&- \Big(\dot{A}_3^2 e^{-2\alpha} - \mathscr{D}\frac{dK}{dI}A_3^2\Big)e^{-2\beta_3} - \Big(\dot{A}_1^2 e^{-2\alpha} - \mathscr{D}\frac{dK}{dI}A_1^2\Big)e^{-2\beta_1}\Big], && \text{(2.20c)}\\
\ddot{\beta}_3 &= -\frac{\kappa e^{2\alpha}}{2}\Big[-\Big(mS + \frac{1}{2}K\frac{d\mathscr{D}}{dS}S - K\mathscr{D}\Big) + \Big(\dot{A}_3^2 e^{-2\alpha} - \mathscr{D}\frac{dK}{dI}A_3^2\Big)e^{-2\beta_3}\\
&- \Big(\dot{A}_1^2 e^{-2\alpha} - \mathscr{D}\frac{dK}{dI}A_1^2\Big)e^{-2\beta_1} - \Big(\dot{A}_2^2 e^{-2\alpha} - \mathscr{D}\frac{dK}{dI}A_2^2\Big)e^{-2\beta_2}\Big]. && \text{(2.20d)}
\end{aligned}
$$

The triviality of off-diagonal components of the Einstein tensor for BI metric leads to

$$T_2^1 = T_3^2 = T_1^3 = 0, \tag{2.21}$$

from which one finds

$$\frac{\dot A_1}{A_1}\frac{\dot A_2}{A_2} = \frac{\dot A_2}{A_2}\frac{\dot A_3}{A_3} = \frac{\dot A_3}{A_3}\frac{\dot A_1}{A_1} = \mathscr{D}\frac{dK}{dI}e^{2\alpha}. \tag{2.22}$$

From (2.22) one easily finds

$$\frac{\dot A_1}{A_1} = \frac{\dot A_2}{A_2} = \frac{\dot A_3}{A_3}, \tag{2.23}$$

leading to the following relations between the three components of vector potential:

$$A_1 = A, \quad A_2 = C_{21}A, \quad A_3 = C_{31}A, \tag{2.24}$$

with $C_{21}$ and $C_{31}$ being constants of integration.

In view of (2.21) and (2.23) one easily finds that

$$\ddot\alpha - \dot\alpha^2 + \dot\beta_1^2 + \dot\beta_2^2 + \dot\beta_3^2 = -\frac{\kappa e^{2\alpha}}{2}\Big[\Big(mS + K\mathscr{D} - \frac{3}{2}K\frac{d\mathscr{D}}{dS}S\Big) - 2\mathscr{D}I\frac{dK}{dI}\Big], \tag{2.25a}$$

$$\ddot\beta_1 = \ddot\beta_2 = \ddot\beta_3 = \frac{\kappa e^{2\alpha}}{2}\Big[mS + \frac{1}{2}K\frac{d\mathscr{D}}{dS}S - K\mathscr{D}\Big]. \tag{2.25b}$$

In view of (2.23) from (2.13) it can be shown that

$$\dot\beta_1 = \dot\beta_2 = \dot\beta_3. \tag{2.26}$$

In view of (2.25b) and (2.26) one concludes that the $\beta$s differ by some constant only, namely

$$\beta_1 = \beta, \quad \beta_2 = \beta + \beta_{21}, \quad \beta_3 = \beta + \beta_{31}, \tag{2.27}$$

with $b$, $\beta_{21}$ and $\beta_{31}$ being some arbitrary constants. Since these constants leads to the different scaling along different axis, there is only one option left, it is to set $\beta_{21} = \beta_{31} = 0$. That means the current model allows isotropization. Inserting $\mathscr{D}\frac{dK}{dI}e^{2\alpha}$ into (2.13) one finds the equation for $A$:

$$A\ddot A + \dot A^2 - 2\dot\beta A\dot A = 0, \tag{2.28}$$

with the solution

$$A = \sqrt{C_1\int e^{2\beta}dt + C_2}, \tag{2.29}$$

where $C_1$ and $C_2$ are arbitrary constants.

On account of (2.24) and (2.27) one now finds

$$I = -QA^2e^{-2\beta}, \quad Q = 1 + C_{21}^2 + C_{31}^2 = \text{const}. \tag{2.30}$$

In view of (2.3), (2.25b) and (2.26) the equation (2.25a) can be rearranged as

$$6\dot\beta^2 = \kappa\Big[2mS - \mathscr{D}\frac{d}{dI}\Big(IK\Big)\Big]e^{2\alpha}. \tag{2.31}$$

In what follows, we consider some some concrete cases.

**Massless spinor field with Heisenberg-Ivanenko nonlinearity**

Let us consider the massless spinor field with Heisenberg-Ivanenko nonlinearity. Note that, in the unified nonlinear spinor theory of Heisenberg, the massive term remains absent, and according to Heisenberg, the particle mass should be obtained as a result of quantization of spinor prematter [20]. In the nonlinear generalization of classical field equations, the massive term does not possess the significance that it possesses in the linear one, as it by no means defines total energy (or mass) of the nonlinear field system. Thus without losing the generality we can consider the massless spinor field putting $m = 0$. For Heisenberg-Ivanenko nonlinearity we have $\mathscr{D}(S) = \sigma S^2$. In this case Eq. (2.25b) takes the form

$$\ddot{\beta} = 0, \tag{2.32}$$

with the solution

$$\beta = bt + b_1, \tag{2.33}$$

where $b$ and $b_1$ are arbitrary constants. In view of (2.33) Eq. (2.34) now reads

$$A\ddot{A} + \dot{A}^2 - 2bA\dot{A} = 0. \tag{2.34}$$

The equation (2.34) allows the following solution:

$$A = \sqrt{e^{2bt} - C}, \tag{2.35}$$

Here $C$ is an arbitrary constant. In what follows, we consider a specific solution to the Eq. (2.34):

$$A = De^{bt}, \tag{2.36}$$

with $D$ being an arbitrary constant. Inserting (2.36) into (2.30) one finds

$$I = -\bar{Q}D^2, \quad \bar{Q} = e^{2b_1}Q = e^{2b_1}[1 + C_{21}^2 + C_{31}^2]. \tag{2.37}$$

On account of (2.36) and $\mathscr{D}$ from the triviality off-diagonal components of energy-momentum tensor we find

$$K = \frac{b^2}{\sigma S_0^2} I + C_3, \tag{2.38}$$

with $C_3$ being some arbitrary constant. Finally inserting $K$ from (2.38) into (2.31) one finds

$$(6 + 2\kappa\bar{Q}D^2)b^2 = \kappa\sigma S_0^2 C_3. \tag{2.39}$$

Equation (2.39) gives the relation between different constants.

**Case with minimal coupling**

Let us consider the case with minimal coupling. In this case from the Off-diagonal components of energy-momentum tensor we find

$$\dot{A}_1\dot{A}_2 = \dot{A}_2\dot{A}_3 = \dot{A}_3\dot{A}_1 = 0. \tag{2.40}$$

From (2.40) follows that at least two of the three components $A_i$ are constant, which means only one of the components of $F_{\mu\nu}$ is nonzero. Let us assume that $\dot{A}_1 = \dot{A} \neq 0$. In view of $\dot{A}_2 = \dot{A}_3 = 0$ from the electromagnetic field equations in this case we have

$$A = C\int e^{2\beta_1}dt + C_1, \quad A_2 = \text{const.}, \quad A_3 = \text{const.}, \tag{2.41}$$

with $C$ and $C_1$ being some arbitrary constants. Components of the energy-momentum tensor in this case read

$$T_0^0 - mS = T_1^1 = -T_2^2 = -T_3^3 = \frac{C_1^2}{2} e^{2\beta_1 - 2\alpha}. \tag{2.42}$$

Einstein field equations in this case takes the form

$$\begin{aligned}
\ddot{\alpha} - \dot{\alpha}^2 + \dot{\beta}_1^2 + \dot{\beta}_2^2 + \dot{\beta}_3^2 &= -\frac{\varkappa C_1^2}{2} e^{2\beta_1} - \frac{m\varkappa S_0}{2} e^{\alpha}, && \text{(2.43a)}\\
\ddot{\beta}_1 &= -\frac{\varkappa C_1^2}{2} e^{2\beta_1} + \frac{m\varkappa S_0}{2} e^{\alpha}, && \text{(2.43b)}\\
\ddot{\beta}_2 &= \frac{\varkappa C_1^2}{2} e^{2\beta_1} + \frac{m\varkappa S_0}{2} e^{\alpha}, && \text{(2.43c)}\\
\ddot{\beta}_3 &= \frac{\varkappa C_1^2}{2} e^{2\beta_1} + \frac{m\varkappa S_0}{2} e^{\alpha}. && \text{(2.43d)}
\end{aligned}$$

Unlike the case with interacting electromagnetic and scalar fields, in case of minimal coupling we have $\ddot{\beta}_1 \neq \ddot{\beta}_2 = \ddot{\beta}_3$, which shows the space-time in this case essentially anisotropic. Addition of (2.43b), (2.43c) and (2.43d), on account of coordinate condition, gives

$$\ddot{\alpha} = \frac{\varkappa C_1^2}{2} e^{2\beta_1} + \frac{3m\varkappa S_0}{2} e^{\alpha}. \tag{2.44}$$

In case of massless spinor field the system (2.43) can be easily solved to obtain

$$e^{2\beta_1} = -\frac{2\eta^2}{\varkappa C_1^2} \mathrm{cosech}^2(\eta t), \quad \eta^2 = \text{const.}, \tag{2.45}$$

and

$$e^{2\alpha} = e^{2\beta_2} = e^{2\beta_3} = \sinh^2(\eta t). \tag{2.46}$$

Thus we see that direct interaction is essential for isotropization process of initially anisotropic space-time.

## III. CONCLUSION

Within the framework of Bianchi type-I cosmological model evolution of the initially anisotropic space-time in presence of an interacting system of spinor and electromagnetic fields is studied. It is shown that the interacting term can be viewed as effective photon mass. The present model allows asymptotic isotropization of initially anisotropic space-time.

---

[1] A.S. Goldhaber and M.M. Nieto, Rev. Mod. Phys. **43**, 277 (1971)

[2] K.D. Froome and L. Essen, *The velocity of light and radio waves* Academis, NY (1969).

[3] B.N. Taylor, H.W. Parker, and D.N. Langenberg, Rev. M od. Phys. **41**, 375 (1969).

[4] L. de Broglie, *Theorie generale des particles a spin* Gauthier-Villars, Paris, 2 nd Ed. (1954).

[5] A.H. Rosenfeld *et al* Rev. Mod. Phys. **40**, 77 (1968).

[6] E.R. Williams, J.E. Faller, and H. Hill, Phys. Rev. Lett. **26**, 721 (1971)

[7] R.E. Crandall, Am. J. Phys. **51**, 698 (1983).
[8] M.A. Chernikov, C.J. Gerber, H.R. Ott, and H.J. Gerber, Phys. Rev. Lett. **68**, 3383 (1992).
[9] B.E. Schaefer, Phys. Rev. Lett. **82**, 4964 (1999).
[10] E. Fishbach *et al* Phys. Rev. Lett. **73**, 514 (1994).
[11] L. Davis, A.S. Goldhaber, and M.M. Nieto, Phys. Rev. Lett. **35**, 1402 (1975).
[12] R. Lakes, Phys. Rev. Lett. **80**, 1826 (1998).
[13] J. Luo, L.-C. Tu, Z.-K. Hu, and E.-J. Luan, Phys. Rev. Lett. **90**, 081801 (2003).
[14] Yu.A. Popov, Yu.P. Rybakov, G.N. Shikin, and Bijan Saha *Electromagnetic field with induced massive term: Case with spinor field* ArXiv:
[15] B. Saha and G.N. Shikin, J. Math. Phys. **38**, 5305 (1997).
[16] Bijan Saha, Phys. Rev. D **64**, 123501 (2001).
[17] Bijan Saha, Physics of Particles and Nuclei **37** Suppl. 1, S13, (2006).
[18] Bijan Saha, Phys. Rev. D **74**, 124030 (2006)
[19] Bijan Saha, Physics of Particles and Nuclei **40**(5),656 (2009).
[20] W. Heisenberg, *Introduction to the unified field theory of elementary particles* (Interscience Publ., London. 1966).